
Introduction to microwave modeling

Motivations for SMRT

Context

Many snow microwave emission models have been developed over the last 30-40 years. The most “generic” ones are widely used by the PM community: HUT, MEMLS, DMRT-QMS, DMRT-ML, ...

A few snow radar backscatter models have been developed by the AM community (side looking radar and nadir altimetry).

- several in specific studies
- DMRT-QMS (L. Tsang's group) is dual mode
- MEMLS has recently been extended to active mode


In this introduction lecture:

- Why such a diversity ?
- Is this diversity apparent or profound ?
- Is this diversity beneficial or counter-productive for the community ?
- What about the dual mode ? Good or bad ?

- Why a new model ?


Microwave model ingredients

- Snowpack
- Ground / sea ice / lake ice
- Forest, Atmosphere, ...



Microwave model ingredients



- Thermal Emission
- Scattering and absorption processes
- Reflection and refraction
- Inter-layer interferences
(e.g. ice crust, L-band, ...)



Microwave model ingredients

Snow is a **dense media** from the perspective of EM wave:

- Scattered by many particles \rightarrow change effective incident field
 - Multiple scattering between particles
- \rightarrow Concept of effective permittivity and Born approximation(s)



- Multi-species (e.g. wet snow)


Microwave model ingredients

Models differ in the ingredients and how detailed is each component described e.g. HUT (Snow + atmosphere) versus DMRT-ML (snow) + RTTOV (atmosphere)

Other constraints:

- frequency range

For typical snowpack:



SMRT is definitely a RT model. The following is about RT models

Microwave model ingredients

Other constraints (cont.):

- Application context → performance, adjoint needed, ...
- language
- license
- ecosystem around the model, documentation, support, training, ...
- collaboration network, institutional constraints, community

There are many good reasons for different models.

But: our community is not so 'big'. Question: are the differences profound or superficial ?

The following is mostly based on « Are existing snow microwave emission models so different ? », Picard et al. AGU 2015

Microwave model ingredients

Radiative transfer models in general:

The radiative transfer equation

$$\mu \frac{\partial \mathbf{I}(\mu, \phi, z)}{\partial z} = -\kappa_e(\mu, \phi, z) \mathbf{I}(\mu, \phi, z) + \frac{1}{4\pi} \iint_{4\pi} \mathbf{P}(\mu, \phi; \mu', \phi', z) \mathbf{I}(\mu', \phi', z) d\Omega' + \kappa_a(\mu, \phi, z) \alpha T(z) \mathbf{1}$$

accompanied with boundary conditions:

$$\begin{aligned} \mathbf{I}^{(l)}(\mu < 0, \phi, z_{l-1}) &= \mathbf{R}^{\text{spec,top},(l)}(\mu) \mathbf{I}^{(l)}(-\mu, \phi, z_{l-1}) + \frac{1}{2\pi} \iint_{2\pi, \mu' > 0} \mathbf{R}^{\text{diff,top},(l)}(\mu, \mu', \phi - \phi') \mathbf{I}^{(l)}(\mu', \phi', z_{l-1}) d\Omega' \\ &+ \mathbf{T}^{\text{spec,bottom},(l-1)}(\mu) \mathbf{I}^{(l-1)}(\mu, \phi, z_{l-1}) + \frac{1}{2\pi} \iint_{2\pi, \mu' < 0} \mathbf{T}^{\text{diff,bottom},(l-1)}(\mu, \mu', \phi - \phi') \mathbf{I}^{(l-1)}(\mu', \phi', z_{l-1}) d\Omega' \end{aligned}$$

Microwave model ingredients

Radiative transfer models in general:

The radiative transfer equation

$$\mu \frac{\partial \mathbf{I}(\mu, \phi, z)}{\partial z} = -\kappa_e(\mu, \phi, z) \mathbf{I}(\mu, \phi, z) + \frac{1}{4\pi} \iint_{4\pi} \mathbf{P}(\mu, \phi; \mu', \phi', z) \mathbf{I}(\mu', \phi', z) d\Omega' + \kappa_a(\mu, \phi, z) \alpha T(z) \mathbf{1}$$

Accompanied with boundary conditions:

$$\begin{aligned} \mathbf{I}^{(l)}(\mu < 0, \phi, z_{l-1}) &= \mathbf{R}^{\text{spec, top}, (l)}(\mu) \mathbf{I}^{(l)}(-\mu, \phi, z_{l-1}) + \frac{1}{2\pi} \iint_{2\pi, \mu' > 0} \mathbf{R}^{\text{diff, top}, (l)}(\mu, \mu', \phi - \phi') \mathbf{I}^{(l)}(\mu', \phi', z_{l-1}) d\Omega' \\ &+ \mathbf{T}^{\text{spec, bottom}, (l-1)}(\mu) \mathbf{I}^{(l-1)}(\mu, \phi, z_{l-1}) + \frac{1}{2\pi} \iint_{2\pi, \mu' < 0} \mathbf{T}^{\text{diff, bottom}, (l-1)}(\mu, \mu', \phi - \phi') \mathbf{I}^{(l-1)}(\mu', \phi', z_{l-1}) d\Omega' \end{aligned}$$


Computation:

- Step 1
- a - compute layer electromagnetic intrinsic properties (κ_e , κ_s , κ_a , \mathbf{P} , ϵ)
 - b - compute interfaces electromagnetic intrinsic properties (\mathbf{R} , \mathbf{T})


Step 2 solve the radiative transfer equation

Comparison of models

It's incredible how different they look:



Step1



Fortran / Matlab

Matlab / Fortran

Matlab

Fortran/Python

FMI

C. Mätzler & co

L. Tsang & co

LGGE (now IGE)

Comparison of models

Numerical comparisons showed that none of the models is significantly/always better than the others.

Tedesco et al. 2006, « Intercomparison of Electromagnetic Models for Passive Microwave Remote Sensing of Snow »

Tian, B. « Quantifying inter-comparison of the microwave emission model of layered snowpacks (MEMLS) and the multilayer dense media radiative transfer theory (DMRT) in modeling snow microwave radiance (IGARSS) », 2010

L. Brucker et al. 2011, thesis and « Modeling time series of microwave brightness temperature at Dome C, Antarctica, using vertically resolved snow temperature and microstructure measurements »

Roy et al. 2013, « Brightness temperature simulations of the Canadian seasonal snowpack driven by measurements of snow specific surface area »

Kwon, Y, « Error Characterization of Coupled Land Surface-Radiative Transfer Models for Snow Microwave Radiance Assimilation », 2015

Roy, A., A. Royer, O. St-Jean-Rondeau, B. Montpetit, G. Picard, A. Mavrovic, N. Marchand, and A. Langlois, Microwave snow emission modeling uncertainties in boreal and subarctic environments, *The Cryosphere* 10, 623-638, doi:10.5194/tc-10-623-2016, 2016

Sandells, M., Essery, R., Rutter, N., Wake, L., Leppänen, L., and Lemmetyinen, J.: Microstructure representation of snow in coupled snowpack and microwave emission models, *The Cryosphere*, 11, 229-246, tc-11-229-2017, 2017


Royer A., A. Roy, B. Montpetit, O. Saint-Jean-Rondeau, G. Picard, L. Brucker, and A. Langlois, Comparison of commonly-used microwave radiative transfer models for snow remote sensing. *Remote Sensing of Environment*, 190, 247—259, doi:10.1016/j.rse.2016.12.020, 2017

Performing a fair comparison is challenging because of the many different components and the different « grain size » metrics (microstructure).

Comparison of models

This talk:

Are existing snow microwave emission models so different?
aren't



Reconcilate:

- the different **electromagnetic theories**
- the different **micro-structure representation** used by these models
- the different **solutions** of the radiative transfer equation

Recent studies: Löwe and Picard (TC, 2015) and Pan et al. (2016)

Guess who ?

Löwe and Picard (TC, 2015)

$$K^2 = k^2 + \frac{f_v(k_s^2 - k^2)}{1 + \frac{k_s^2 - k^2}{3K^2}(1 - f_v)} \left\{ 1 + i \frac{2(k_s^2 - k^2)Ka^3}{9 \left[1 + \frac{(k_s^2 - k^2)}{3K^2}(1 - f_v) \right]} \times \left[1 + 4\pi n \int_0^\infty dr r^2 (g(r) - 1) \right] \right\}$$

$$\tilde{\omega} = \frac{2a^3 f_v}{9 \kappa_e} \left| \frac{k_s^2 - k^2}{1 + \frac{k_s^2 - k^2}{3K^2}(1 - f_v)} \right|^2 \left[1 + 4\pi n \int_0^\infty dr r^2 (g(r) - 1) \right]$$

$$\gamma^{\text{bi}}(\hat{\mathbf{o}}, \hat{\mathbf{i}}) = \frac{k_{\text{eff}}^4}{4\pi V} |\mathbf{F}_f|^2 \sin^2 \chi$$

$$= \nu(1 - \nu)(\epsilon_2 - \epsilon_1)^2 K^2 I \cdot k^4 \sin^2 \chi \quad I = \frac{1}{\alpha} \int_0^\infty A(x) x \sin(\alpha x) dx$$

$$\epsilon_{\text{eff}} = \frac{2\epsilon_1 - \epsilon_2 + 3\nu(\epsilon_2 - \epsilon_1) + \sqrt{(2\epsilon_1 - \epsilon_2 + 3\nu(\epsilon_2 - \epsilon_1))^2 + 8\epsilon_1\epsilon_2}}{4}$$

“IBA” and “DMRT QCA-CP” theories in MEMLS and DMRT*

Löwe and Picard, 2015, in the low frequency limit (≤ 37 GHz for most snow) for spherical scatters

- Effective medium permittivity/wavenumber:

$$\varepsilon_{\text{eff},0}^{\text{IBA}} = \frac{2\varepsilon_1 - \varepsilon_2 + 3\phi_2(\varepsilon_2 - \varepsilon_1)}{4} + \frac{\sqrt{(2\varepsilon_1 - \varepsilon_2 + 3\phi_2(\varepsilon_2 - \varepsilon_1))^2 + 8\varepsilon_1\varepsilon_2}}{4}$$


$$\varepsilon_{\text{eff},0}^{\text{QCA-CP}} = \frac{\varepsilon_1 - \frac{(\varepsilon_2 - \varepsilon_1)}{3}(1 - 4\phi_2)}{2} + \frac{\sqrt{\left(\varepsilon_1 - \frac{(\varepsilon_2 - \varepsilon_1)}{3}(1 - 4\phi_2)\right)^2 + 4\varepsilon_1 \frac{(\varepsilon_2 - \varepsilon_1)}{3}(1 - \phi_2)}}{2}$$

- Absorption formulations are identical

- Scattering coefficients:

$$\kappa_s^{\text{IBA}} = \frac{2}{9}k_0^4 a^3 \phi_2 \left| \frac{(\varepsilon_2 - \varepsilon_1) \left(2\varepsilon_{\text{eff},0}^{\text{IBA}} + \varepsilon_1\right)}{\left(2\varepsilon_{\text{eff},0}^{\text{IBA}} + \varepsilon_2\right)} \right|^2 S(0)$$

$$\kappa_s^{\text{QCA-CP}} = \frac{2}{9}k_0^4 a^3 \phi_2 \left| \frac{3\varepsilon_{\text{eff},0}^{\text{QCA-CP}} (\varepsilon_2 - \varepsilon_1)}{3\varepsilon_{\text{eff},0}^{\text{QCA-CP}} + (\varepsilon_2 - \varepsilon_1)(1 - \phi_2)} \right|^2 S(0).$$





Micro-structure in the models

$S(0)$ = snow micro-structure

DMRT-QCA

IBA

Position of the scatterers



Tied to the concept of scatterers. Shape, size and position are not coupled

Micro-structure can be any discrete biphasic medium

Pair-correlation $\mathbf{g}(\mathbf{r}) \sim$ Probability of distance between centres of the scatterers

Autocorrelation of the indicator function $\mathbf{C}(\mathbf{r}) \sim$ Probability of the distance between masses

Distribution

Sticky hard sphere

Exponential autocorrelation function


Parameters

Radius, stickiness

Correlation length (p_{ex})

Micro-structure in the models

When IBA uses Sticky Hard Sphere like DMRT instead of exponential autocorrelation:



Conclusion:

The main difference between MEMLS and DMRT family is the microstructure

Micro-structure in the models

HUT has semi-empirical formulation of scattering/extinction coefficient

Grain size d_0

$$\kappa_e = 0.0018 f^{2.8} d_0^{2.0} \quad \text{where } d_0 = 1.5 (1 - \exp(-1.5 D_{\max}))$$


$$\kappa_s^{\text{IBA}} = \frac{2}{9} k_0^4 a^3 \phi_2 \left| \frac{(\varepsilon_2 - \varepsilon_1) (2\varepsilon_{\text{eff},0}^{\text{IBA}} + \varepsilon_1)}{(2\varepsilon_{\text{eff},0}^{\text{IBA}} + \varepsilon_2)} \right|^2 S(0)$$

$$\kappa_s^{\text{QCA-CP}} = \frac{2}{9} k_0^4 a^3 \phi_2 \left| \frac{3\varepsilon_{\text{eff},0}^{\text{QCA-CP}} (\varepsilon_2 - \varepsilon_1)}{3\varepsilon_{\text{eff},0}^{\text{QCA-CP}} + (\varepsilon_2 - \varepsilon_1) (1 - \phi_2)} \right|^2 S(0).$$

Using micro-structure images \rightarrow a geometrically-based relationship between D_{\max} and p_{ex}

Micro-structure in the models

Pan, Durand and co-authors, 2016




HUT and IBA have very different scattering coefficients !


Surprising because HUT and MEMLS are known to have good performance...

Micro-structure in the models

HUT : snow is a strongly forward scattering : $q=0.96$



IBA and QCA-CP : snow scattering is almost isotropic (Rayleigh or moderate Mie)



Similarity theory

Radiative transfer equation:

$$\mu \frac{\partial \mathbf{I}(\mu, \phi, z)}{\partial z} = -\kappa_e(\mu, \phi, z) \cdot \mathbf{I}(\mu, \phi, z) + \frac{1}{4\pi} \iint_{4\pi} \mathbf{P}(\mu, \phi; \mu', \phi', z) \cdot \mathbf{I}(\mu', \phi', z) d\Omega' + \kappa_a(\mu, \phi, z) T(z)$$

Different formulations of κ_e and \mathbf{P} may lead to exactly the same RT equation (and exactly the same solution)

e.g.

C. Mitrescu, , G.L. Stephens, On similarity and scaling of the radiative transfer equation, Journal of Quantitative Spectroscopy and Radiative Transfer 86, 4, 387–394, 2004

H.C. van de Hulst, Multiple light scattering, Academic Press, New York, 1980



Joseph, Wiscombe, Weiman. The Delta-Eddington Approximation for Radiative Flux Transfer. Journal of the Atmospheric Sciences, 1976, 33, 2452-2459.

Similarity theory

Visible in the two-flux theory: single scattering albedo ω ($\sim K_s$) and asymmetry factor g ($\sim P$):

$$\omega, g \leftrightarrow \omega', g' \quad \omega' = \frac{(1-f)\omega}{1-f\omega} \quad g' = \frac{g-f}{1-f} \quad \text{for any } f$$


M-delta approximation: choose f to reduce the forward peak ($0 < f < g$)



Mie calculation

Similarity theory

Pan, Durand and co-authors, 2016



... good agreement once the comparison takes into account the different phase function shape → **HUT has similar behavior as MEMLS despite huge apparent differences**

Conclusion

Back to the introductory questions, my opinionated response:

- Is this diversity apparent or profound ?

- overall apparent
- all models converge to the “right” snow behaviour and give reasonable results (not always for physically correct reasons)

- Why such a diversity ?

- historical
- different focus/approach

- Is this diversity beneficial or counter-productive for the community ?

- it has been beneficial until many users started to be spend more time performing numerical inter-comparisons (incl. myself) than really using models to develop useful algorithms for end-users.

- What about the dual mode ? Good or bad ?

- it's time to merge both because of dual mode missions and in-situ datasets

- Why a new model ?

Conclusion

- Why a new model ?

We don't need a new model (yet) but we need:

a repository of microwave community knowledge

= merge all RT models / theories in one code base, one framework

with extended capabilities to explore the micro-structure

with multi mode capabilities (passive, radar, altimeter)

with easier access for beginners and non-specialists

using modern and more efficient languages and programming techniques

Conclusion

